



PubMed	Nucleotide	Protein	Genome	Structure	PMC	Taxonomy	OMIM	Books		
Search	PubMed	<input type="text"/>	for	<input type="text"/>	Go	Clear				
Limits    Preview/Index    History    Clipboard    Details										
Display	Abstract	<input type="text"/>	Show:	20	<input type="text"/>	Sort	<input type="text"/>	Send to	Text	<input type="text"/>

Entrez  
PubMed☐ 1: J Clin Microbiol 1997 Mar;35(3):570-7

Related Articles, Links

FREE full text article at  
[jcm.asm.org](http://jcm.asm.org)

**A one-tube method of reverse transcription-PCR to efficiently amplify a 3-kilobase region from the RNA polymerase gene to the poly(A) tail of small round-structured viruses (Norwalk-like viruses).**

PubMed  
Services**Ando T, Monroe SS, Noel JS, Glass RI.**

Viral Gastroenteritis Section, National Center for Infectious Diseases, Centers for Disease Control and Prevention, Atlanta, Georgia 30333, USA.  
[txa5@CIDDDVD1.em.cdc.gov](mailto:txa5@CIDDDVD1.em.cdc.gov)

Related  
Resources

Amplification of a 3-kb genome region from the RNA polymerase gene to the 3' poly(A) tail of small round-structured virus (SRSV) by reverse transcription-PCR (RT-PCR) has been difficult to achieve because of a stable secondary structure in a region between the RNA polymerase gene and the 5' end of the second open reading frame. We have developed a one-tube RT-PCR method to efficiently amplify this region. The method comprises three procedures: purification of poly(A)+ RNA from a starting RNA solution by oligo(dT)30 covalently linked to latex particles, buffer exchange, and continuous RT and PCR in a single tube containing all reaction components. The key elements of this method are (i) first-strand cDNA synthesis with the Superscript II version of RNase H- Moloney murine leukemia virus reverse transcriptase at 50 degrees C for 10 min by using the RNA-oligo(dT)30 hybrid on the latex particles as the template and primer, and (ii) PCR by Taq and Pwo DNA polymerases mixed together with a mixture of 12 phased oligo(dT)25 antisense primers. The detection threshold of the one-tube RT-PCR method was as little as 0.2 ng of the crude RNA used as the source of the template. Using this method, we obtained 3-kb products from 24 SRSV strains previously characterized into four genetic groups. These included 5 P1-A, 4 P1-B, 5 P2-A, and 10 P2-B strains. Because SRSVs have not yet been cultivated in vitro, this novel method should facilitate molecular characterization of SRSVs to provide a firm scientific foundation for improvements and refinements of SRSV diagnostics.


PMID: 9041391 [PubMed - indexed for MEDLINE]

Display	Abstract	▼	Show:	20	▼	Sort	▼	Send to	Text	▼
---------	----------	---	-------	----	---	------	---	---------	------	---

[Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)  
[Department of Health & Human Services](#)  
[Freedom of Information Act](#) | [Disclaimer](#)

Feb 19 2003 14:38:47

■



PubMed	Nucleotide	Protein	Genome	Structure	PMC	Taxonomy	OMIM	Books
Search PubMed	▼ for longRT-PCR using Pwo and taq DNA polymerases						Go	Clear
Limits		Preview/Index		History		Clipboard		Details

One of your terms is not found in the database.

Entrez  
PubMed

Display	Summary	▼	Show: 20	▼	Sort	▼	Send to	Text	▼
Items 1-14 of 14								One page.	

PubMed  
Services

- ☐ **1:** [Held HA, Benner SA.](#) Related Articles, Links  
Challenging artificial genetic systems: thymidine analogs with 5-position sulfur functionality.  
Nucleic Acids Res. 2002 Sep 1;30(17):3857-69.  
PMID: 12202771 [PubMed - indexed for MEDLINE]

- ☐ **2:** [Attatippaholkun WH, Attatippaholkun MK, Nisalak A, Vaughn DW, Innis BL.](#) Related Articles, Links  
A novel method for the preparation of large cDNA fragments from dengue-3 RNA genome by long RT-PCR amplification.  
Southeast Asian J Trop Med Public Health. 2000;31 Suppl 1:126-33.  
PMID: 11414441 [PubMed - indexed for MEDLINE]

Related  
Resources

- ☐ **3:** [Mullan B, Kenny-Walsh E, Collins JK, Shanahan F, Fanning LJ.](#) Related Articles, Links  
Inferred hepatitis C virus quasispecies diversity is influenced by choice of DNA polymerase in reverse transcriptase-polymerase chain reactions.  
Anal Biochem. 2001 Feb 15;289(2):137-46.  
PMID: 11161307 [PubMed - indexed for MEDLINE]

- ☐ **4:** [Forrest R, Zhou H, Hickford JG.](#) Related Articles, Links  
3'overhangs influence PCR-SSCP patterns.  
Biotechniques. 2000 Nov;29(5):958-60, 962. No abstract available.  
PMID: 11084854 [PubMed - indexed for MEDLINE]

- ☐ **5:** [Kainz P.](#) Related Articles, Links  
The PCR plateau phase - towards an understanding of its limitations.  
Biochim Biophys Acta. 2000 Nov 15;1494(1-2):23-7.  
PMID: 11072065 [PubMed - indexed for MEDLINE]

- ☐ **6:** [Fromenty B, Demeilliers C, Mansouri A, Pessayre D.](#) Related Articles, Links  
Escherichia coli exonuclease III enhances long PCR amplification of damaged DNA templates.  
Nucleic Acids Res. 2000 Jun 1;28(11):E50.  
PMID: 10871349 [PubMed - indexed for MEDLINE]

- ☐ **7:** [Schulze E, Bettendorf M, Maser-Gluth C, Decker M, Schwabe U.](#) Related Articles, Links  
Allele-dropout using PCR-based diagnosis for the splicing mutation in intron-2 of

the CYP21B-gene: successful amplification with a Taq/Pwo-polymerase mixture.  
Endocr Res. 1998 Aug-Nov;24(3-4):637-41.  
PMID: 9888552 [PubMed - indexed for MEDLINE]

☐ 8: [Liu Q, Sommer SS.](#) Related Articles, Links

Subcycling-PCR for multiplex long-distance amplification of regions with high and low GC content: application to the inversion hotspot in the factor VIII gene.  
Biotechniques. 1998 Dec;25(6):1022-8.  
PMID: 9863056 [PubMed - indexed for MEDLINE]

☐ 9: [Iannelli F, Giunti L, Pozzi G.](#) Related Articles, Links

Direct sequencing of long polymerase chain reaction fragments.  
Mol Biotechnol. 1998 Oct;10(2):183-5.  
PMID: 9819816 [PubMed - indexed for MEDLINE]

☐ 10: [Abu Al-Soud W, Radstrom P.](#) Related Articles, Links

Capacity of nine thermostable DNA polymerases To mediate DNA amplification in the presence of PCR-inhibiting samples.  
Appl Environ Microbiol. 1998 Oct;64(10):3748-53.  
PMID: 9758794 [PubMed - indexed for MEDLINE]

☐ 11: [Gunther S, Sommer G, Von Breunig F, Iwanska A, Kalinina T, Sterneck M, Will H.](#) Related Articles, Links

Amplification of full-length hepatitis B virus genomes from samples from patients with low levels of viremia: frequency and functional consequences of PCR-introduced mutations.  
J Clin Microbiol. 1998 Feb;36(2):531-8.  
PMID: 9466771 [PubMed - indexed for MEDLINE]

☐ 12: [Rispeter K, Lu M, Lechner S, Zibert A, Roggendorf M.](#) Related Articles, Links

Cloning and characterization of a complete open reading frame of the hepatitis C virus genome in only two cDNA fragments.  
J Gen Virol. 1997 Nov;78 ( Pt 11):2751-9.  
PMID: 9367360 [PubMed - indexed for MEDLINE]

☐ 13: [Ando T, Monroe SS, Noel JS, Glass RI.](#) Related Articles, Links

A one-tube method of reverse transcription-PCR to efficiently amplify a 3-kilobase region from the RNA polymerase gene to the poly(A) tail of small round-structured viruses (Norwalk-like viruses).  
J Clin Microbiol. 1997 Mar;35(3):570-7.  
PMID: 9041391 [PubMed - indexed for MEDLINE]

☐ 14: [Maruyama H, Kawakami H, Nakamura S.](#) Related Articles, Links

Reevaluation of the exact CAG repeat length in hereditary cerebellar ataxias using highly denaturing conditions and long PCR.  
Hum Genet. 1996 May;97(5):591-5.  
PMID: 8655136 [PubMed - indexed for MEDLINE]

Display	Summary	▼	Show: 20	▼	Sort	▼	Send to	Text	▼
Items 1-14 of 14								One page.	

[Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)  
[Department of Health & Human Services](#)  
[Freedom of Information Act](#) | [Disclaimer](#)

Feb 19 2003 14:38:47

■